



HOGER DAROO

Pharmaceutical Company

Mirtager® ftilm-coated tablets

Dosage form and active ingredients:

Each film coated circle tablet contains 30 mg of Mirtazapine.

Therapeutic indications

Mirtazapine is one of a group of medicines called antidepressants and is used to treat depressive illness.

Drug Dosage and Administration method:

The usual starting dose is 15 or 30 mg every day.

Your doctor may advise you to increase your dose after a few days to the amount that is best for you (between 15 and 45 mg per day).

Treatment with Mirtager® should only be initiated by physicians.

Medicine taking time:

Take Mirtager® tablets at a specific time in a day or before sleeping.

Contraindications:

● In the case of allergy (severe allergic reaction) to Mirtager® or other supplements of this drug

- Recent history of the use of monoamine oxidase inhibitors (MOA-Is)
- Having suicidal thoughts
- Mirtager® pills are not usually used in children and adolescents below 18 years of age due to their unclear profile.

Pregnancy and lactation:

- Talk to your doctor before taking any medicine
- Limited experience of Mirtager® use in pregnant women does not indicate an increased risk, however, during pregnancy caution should be exercised.
- If you have taken migraine tablets before or shortly before your birth, you should check your baby for possible side effects.
- Your breastfeeding will be prescribed by your doctor during taking Mirtager® pills.

Driving and working with machinery:

Mirtager® pills can affect your concentration and vigilance

Side effects:

- Increase appetite and weight
- Drowsiness
- Headache
- Dry mouth

If any of the following side effects occur while taking Mirtager®, check with your doctor.

- Symptoms of infection such as fever, sore throat, mouth ulcers.
- Feeling great happiness and excitement (Mania)

- Yellow skin and eyes (jaundice symptoms)
- Epilepsy (seizures)
- Severe fever, sweating, increased heart rate, diarrhea, muscle contractions (uncontrollable), tremor, involuntary hyperactivity reactions, restlessness, mood changes and unconsciousness.
- Any thoughts about suicide or self-harm while taking this medicine.

Storage condition:

- Store bellow 30 degree centigrade and keep away from direct light, moist and frizzing.
- Keep out of reach of children.
- Don't use expired Mirtager®.

Nature and contents of container

10 Tablets are packaged in each opaque blister cards and each package contains 30 tablets.

Marketing authorization holder

Hogerdaroo danesh Phtarmaceutical CO.
Shahid Beheshti Uni. Technology Incubators
Tehran, Iran.

Postal Code: 1983969411

Packing site:

Kishmedipham Pharmaceutical Company-
kish island, iran.



HOGER DAROO

Pharmaceutical Company



KISHMEDIPHARM

Caring your health

www.kishmedipham.com